

Cuadro 3. Contenido de ácidos grasos en aceite de semilla de genotipos de nopal (*Opuntia* spp.) sobresalientes en producción de fruto.

Genotipo	Contenido (%)							
	Palmitico	Estearico	Oleico	Linoleico	linolenico	Sat/Insat	Oleic/ Linoleico	Linoleico/ Linolenico
1	11.89 ± 0.11	2.75 ± 0.45	11.32 ± 0.21	69.52 ± 0.25	4.52 ± 0.12	0.17 ± 0.00	0.16 ± 0.00	15.38 ± 0.46
2	13.23 ± 0.19	3.79 ± 0.17	11.38 ± 0.43	66.75 ± 0.19	4.85 ± 0.12	0.21 ± 0.01	0.17 ± 0.01	13.76 ± 0.38
3	11.81 ± 0.01	3.79 ± 0.03	10.38 ± 0.04	70.32 ± 0.00	3.70 ± 0.00	0.18 ± 0.00	0.15 ± 0.00	19.02 ± 0.02
4	11.11 ± 0.05	2.88 ± 0.66	11.48 ± 1.17	67.57 ± 0.91	4.52 ± 0.76	0.17 ± 0.00	0.17 ± 0.02	15.14 ± 2.34
5	13.23 ± 0.17	2.82 ± 0.14	14.46 ± 0.30	63.90 ± 0.04	5.59 ± 0.06	0.19 ± 0.00	0.23 ± 0.00	11.44 ± 0.12
6	12.59 ± 0.15	2.92 ± 0.52	9.75 ± 0.25	69.38 ± 0.75	5.37 ± 0.17	0.18 ± 0.01	0.14 ± 0.01	12.93 ± 0.27
7	11.90 ± 0.20	2.72 ± 0.01	16.75 ± 0.14	64 ± 0.27	4.64 ± 0.07	0.17 ± 0.00	0.26 ± 0.00	13.80 ± 0.15
8	11.51 ± 0.50	2.28 ± 0.11	11.12 ± 0.14	69.74 ± 0.58	5.35 ± 0.04	0.16 ± 0.01	0.16 ± 0.00	13.03 ± 0.20
9	14.28 ± 0.06	3.21 ± 0.29	12.22 ± 0.08	65.10 ± 0.02	5.19 ± 0.13	0.21 ± 0.00	0.19 ± 0.00	12.56 ± 0.30
10	11.37 ± 0.01	2.61 ± 0.12	13.20 ± 0.14	63.24 ± 7.35	4.57 ± 0.00	0.17 ± 0.02	0.21 ± 0.03	13.83 ± 1.62
11	10.97 ± 0.03	2.63 ± 0.07	9.98 ± 0.54	73.09 ± 0.56	3.34 ± 0.12	0.16 ± 0.00	0.14 ± 0.01	92.0 ± 0.95
	12.17	2.95	11.19	67.59	4.69	0.18	0.18	14.83